



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Sodium pump regulation of locomotor control circuits

Citation for published version:

Picton, L, Zhang, H & Sillar, KT 2017, 'Sodium pump regulation of locomotor control circuits', *Journal of Neurophysiology*. <https://doi.org/10.1152/jn.00066.2017>

Digital Object Identifier (DOI):

[10.1152/jn.00066.2017](https://doi.org/10.1152/jn.00066.2017)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Journal of Neurophysiology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



1 Sodium pump regulation of locomotor control circuits

2

3 Laurence D Picton, HongYan Zhang* and Keith T Sillar

4

5 School of Psychology and Neuroscience
6 St Mary's Quad
7 University of St. Andrews
8 St Andrews, Fife KY16 9JP, Scotland, UK.

9 *Centre for Neuroregeneration
10 University of Edinburgh
11 The Chancellor's Building
12 49 Little France Crescent, Edinburgh EH16 4SB, Scotland, UK.

13 Acknowledgments:

14 We are grateful for the financial support of the BBSRC (grant numbers:
15 BB/M024946/1 and BB/JO1446X/1), the Carnegie Trust and the University of St
16 Andrews. We also thank Dr Stefan Pulver (School of Psychology and Neuroscience,
17 University of St Andrews) for providing images and data on *Drosophila* third instar
18 larval crawling.

19 Abstract

20 Sodium pumps are ubiquitously expressed membrane proteins that extrude three
21 Na⁺ ions in exchange for two K⁺ ions using ATP as an energy source. Recent studies
22 have illuminated additional, dynamic roles for sodium pumps in regulating the
23 excitability of neuronal networks in an activity-dependent fashion. Here we review
24 their role in a novel form of short-term memory within rhythmic locomotor networks.
25 The data we review derives mainly from recent studies on *Xenopus* tadpoles and
26 neonatal mice. The role and underlying mechanisms of pump action broadly match
27 previously published data from an invertebrate, the *Drosophila* larva. We therefore
28 propose a highly conserved mechanism by which sodium pump activity increases
29 following a bout of locomotion. This results in an ultraslow afterhyperpolarisation
30 (usAHP) of the membrane potential that lasts around 1 minute, but which only occurs
31 in around half the network neurons. This usAHP in turn alters network excitability so
32 that network output is reduced in a locomotor interval-dependent manner. The
33 pumps therefore confer on spinal locomotor networks a temporary memory trace of
34 recent network performance.

35 Introduction

36 Motor systems have evolved to meet the species-specific behavioural requirements
37 upon which animal survival and reproduction depend. To succeed, the underlying
38 motor circuits must be adaptable in the face of the demands placed on individuals by
39 prevailing external and internal conditions. Such circuit adaptations, which may
40 relate to developmental stage and/or hormonal state, are mostly due to changes in
41 the integrative electrical properties of, and synaptic weightings between, component

neurons within motor circuits (Harris-Warrick and Marder 1991). Many of these changes are mediated by the opening of ion channels, and the consequent alterations to circuit function can involve both neuromodulation and activity-dependent neuronal plasticity. One disadvantage of this ion channel-based strategy is that the decrease in input resistance that accompanies channel opening could shunt incoming synaptic inputs and decrease the responsiveness of neurons and subsequent network output. This, in turn, could compromise the intended behaviour, and if this involves the escape from a predator, for example, it could be potentially catastrophic for survival. An alternative strategy is for neuronal activity or neuromodulation to affect the function of ion pumps which, since there is no change in input resistance, should not shunt the membrane response and hence preserve the responsiveness of the network to various inputs. Furthermore, changes in the activity of ion pumps can exert effects on the excitability of neurons on a much slower timescale, over many seconds and even minutes, leaving a prolonged memory trace of a neuron's recent activity.

The $\text{Na}^+\text{-K}^+$ ATPase (*aka* the Na^+ pump) is one of the most ubiquitously expressed proteins in the animal kingdom, which is most renowned for its role in establishing a gradient of high extracellular Na^+ and high intracellular K^+ ion concentrations across cell membranes. With each Na^+ pump cycle, three Na^+ ions are extruded and two K^+ ions flow into the cell, utilizing ATP as an energy source. Because of this charge asymmetry, Na^+ pump activity sets and homeostatically maintains the resting membrane potential upon which neuronal firing relies, and in so doing accounts for more than half of all brain energy consumption (Engl and Attwell 2015).

Recently, a novel and dynamic role for the Na^+ pump as an activity-dependent regulator of brain and spinal circuit function has been reported across a wide range of neurons, systems, behaviours and species. Within motor systems, for example, seminal work on crawling in *Drosophila* larvae has demonstrated that high frequency action potential firing of motoneurons causes a pump-mediated hyperpolarization lasting tens of seconds, which in turn influences future locomotory crawling behaviour (Pulver and Griffith 2010). In the present paper, we review and compare similar findings from spinal central pattern generator (CPG) circuits controlling rhythmic locomotion in two phylogenetically disparate vertebrate model systems: the *Xenopus* frog tadpole and the neonatal mouse. As in *Drosophila*, these circuits also possess an intrinsic pump-based mechanism that links future to past network activity. This suggests a highly conserved, pump-mediated dynamic regulation of motor circuit function. In spinal motor circuits, the duration of a bout of locomotion is influenced by previous network activity if two bouts occur within about a minute of each other; a form of short-term motor memory (Picton et al. 2017; Zhang and Sillar 2012; Zhang et al. 2015). This motor memory relies on the presence of a pump-mediated ultraslow afterhyperpolarization (usAHP) of up to 10 mV in spinal neurons, which lasts for the same duration of approximately a minute.

Na^+ pump regulation in three locomotor systems

The ultra-slow afterhyperpolarisation (the usAHP)

85 In both the tadpole (Figure 1A) and neonatal mouse (Figure 1B), high frequency
86 action potential firing drives the resting membrane potential to a more hyperpolarized
87 level in a subset of motoneurons and interneurons in the spinal cord. A remarkably
88 similar phenomenon has also been reported in *Drosophila* larva motoneurons
89 (Pulver and Griffith 2010; Figure 1C). This hyperpolarization is distinguished from
90 other ion channel-mediated AHPs (e.g. the “fast”, “medium” or “slow” AHP; Storm
91 1987) largely by its duration, with neurons remaining hyperpolarised once activity
92 has stopped for up to one minute. Although the amplitude of a usAHP can vary quite
93 considerably both within and between neuron types, our findings in *Xenopus* and
94 mouse spinal neurons suggest that, on average, the pump AHP involves a
95 hyperpolarization of approximately 5 mV (Figure 1Aii,Bii), remarkably similar to the
96 equivalent event in *Drosophila* larvae (Figure 1Cii).




Figure 1. The ultraslow afterhyperpolarisation (usAHP) in CPG neurons of three species. **Ai.** Experimental preparation for making patch-clamp recordings from an immobilised stage 37/8 *Xenopus* tadpole. **Aii.** Following either swimming, or in this case a long suprathreshold current pulse, the membrane potential is driven to a more hyperpolarised membrane potential (the usAHP). **Bi.** Experimental preparation for making patch-clamp recordings from neonatal mice. **Bii.** Following a long suprathreshold current pulse, a usAHP is observed in spinal motoneurons and interneurons in neonatal mice. **Ci.** Schematic of a third instar *Drosophila* larva. **Cii.** A usAHP observed in a *Drosophila* motoneuron.

Besides its long duration, several other features of the usAHP distinguish it from ion channel-mediated AHP mechanisms. For example, because it is mediated by the Na^+ pump, it is selectively blocked by a low concentration of the cardiac glycoside ouabain (Figure 2Ai,Bi,Ci). The usAHP is also highly dependent on the accumulation of intracellular sodium that accompanies repetitive action potential firing. Therefore blocking fast sodium channels with TTX, to prevent action potential generation, also effectively abolishes the usAHP (Figure 2Aii,Bii,Cii). Thirdly, because the usAHP

occurs upon the increased activation of ion pumps, rather than ion channel opening or closing, there are no detectable changes in conductance, and this can be observed by measuring a consistent membrane response to small injections of hyperpolarising current throughout the usAHP (Figure 2Aii,Bii,Cii). Perhaps not surprisingly, there are a number of differences in the features of the usAHP in tadpoles and mice at the single-cell level. For example, whilst ouabain and TTX completely abolish the usAHP in tadpoles, a shorter-duration AHP often persists in many motoneurons and interneurons in mice (Figure 2Bi,ii), presumably due to the presence of additional, voltage-dependent AHP mechanisms such as the medium and/or slow AHP, which can persist in the absence of spiking (Rekling et al. 2000).




Figure 2. A cross-species comparison of the basic features of the usAHP. **Ai.** The usAHP is abolished by the Na^+ pump blocker ouabain. **Aii.** The usAHP is also abolished when fast Na^+ channels are blocked using TTX. **Aiii.** By measuring the membrane response to small hyperpolarising current pulse we found no changes in conductance before, during or after the induction of a usAHP, suggesting the involvement of a Na^+ pump (adapted from Zhang and Sillar 2012). The experimental manipulations outlined in **A** have similar results in neonatal mouse CPG neurons (**B**; adapted from Picton et al, 2017) and *Drosophila* motoneurons (**C**; adapted from Pulver and Griffith 2010).

Physiological roles for the Na^+ pump

By its very nature, the usAHP is ideally positioned to function as a spike rate monitor, whose duration and amplitude reflects the integration of spike frequency over time. Furthermore, the usAHP is not only generated in response to artificial current injection protocols used to evoke spikes, but by any stimulus that produces trains of action potentials sufficient to generate a build-up of intracellular sodium (e.g. locomotion, Figure 3Aii). Importantly, because the usAHP recovers over a period of around a minute, it acts as a transient engram of how recently, and how intensely locomotor activity occurred.

In *Xenopus* tadpoles, we have explored how this short-term memory of recent activity acts to regulate the interval relationship between evoked episodes of “fictive swimming” (motor output without muscle contraction). When the interval between swim episodes is set to longer than the duration of a usAHP (longer than 1 minute), episodes of evoked swimming in a “well rested” tadpole are statistically identical, both in the duration of a swim episode and all other parameters of swimming (swim frequency, burst durations etc.). However, when this interval is reduced to 30, 15 or 5 seconds, the second episode is progressively shorter, slower and weaker, in an interval-dependent manner (Figure 3B; note spike failures in episode 2, Figure 3C). The importance of the Na⁺ pump for this self-regulation of network output becomes clear when the pumps are blocked by ouabain; the animal becomes completely unable to regulate its own locomotor activity, causing it to swim almost indefinitely (Figure 2D).

The swim durations and inter-episode intervals involved here may seem short anthropomorphically (tens of seconds), but need to be scaled to be appreciated from a human perspective, and in the broader context of locomotion. If we treat a single tail undulation as equivalent to one human stride, then a typical 2 minute episode of 20 Hz swimming (~2400 swim cycles) could be considered broadly equivalent to a 5 km sprint for a human (assuming a typical stride length of ~2 metres). This distance could comfortably be covered in around 30 minutes, but imagine resting only for a minute before being stimulated to sprint again while still fatigued; the runner is unlikely to get as far, or locomote at the same speed, as it could from a well-rested start. Whether Na⁺ pumps play a direct role in human fatigue is not yet completely clear, but certainly the evidence for central mechanisms of fatigue is extremely compelling (reviewed in Gandevia 2001). More specifically, there is strong evidence that central fatigue involves an activity-dependent reduction in motoneuron drive (Ranieri and Di Lazzaro 2012; Rossi et al. 2012). Furthermore, it has been shown that human motor axons display an activity-dependent hyperpolarisation following natural activity, which is due to an enhancement of Na⁺ pump activity, and whose duration and amplitude depends on the axonal discharge rate (Kiernan et al. 2004; Vagg et al. 1998). This raises the fascinating possibility that an activity-dependent enhancement of Na⁺ pump activity in spinal neurons may contribute to fatigue during human locomotion. Given the ubiquity of pumps throughout the nervous system they have enormous potential as drug targets, with important implications not only for endurance athletes, but also in the context of diseases associated with fatigue symptoms such as diabetes (Krishnan et al. 2008) and ALS (Ellis et al. 2003), in which sodium pump dysfunction has been implicated.

It has long been known that one way to experimentally “fatigue” a neuron is to raise the levels of intracellular sodium. These experiments were first conducted on the squid giant axon in the mid 1950’s and, quite unexpectedly, high sodium resulted in a tonic membrane hyperpolarisation (Hodgkin and Keynes 1956) that turned out to be mediated by enhanced Na^+ pump activity. In our experiments, we have used a drug called monensin, a sodium ionophore, to raise the level of intracellular sodium in spinal CPG neurons. This not only enhances the usAHP by increasing Na^+ pump activity, but in effect it causes the locomotor network to become chronically fatigued. Under these conditions, the swim network acts as if it is being activated from an unrested starting point, resulting in weaker, slower and shorter locomotion.




Figure 3. The usAHP as a short-term memory mechanism in *Xenopus* tadpoles. **Ai.** Schematic showing the experimental set-up. **Aii.** A brief (1 ms) current pulse to the tail (Stim.) initiates an episode of swimming which is recorded at both the single cell level (Aii, top) and at the level of overall network output using ventral root recording (Aii, bottom). Note the prolonged membrane hyperpolarisation (usAHP) in the intracellular trace at the end of the swim episode. Inset shows an expansion of the recording indicated by the black box showing the intracellular and ventral root traces during swimming. **B.** Ventral root recordings showing that an evoked swim episode is shorter and slower when it follows a previous episode after a 5, 15 or 30 second interval. **C.** The interval relationship is apparent when activity is evoked within the 1 minute usAHP that follows swimming, which reduces the spike probability of CPG neurons. **D.** Real swimming behaviour in a *Xenopus* tadpole with multiple consecutive video frames overlapped to show swim path in response to touch. When the Na^+ pumps are blocked using ouabain the tadpole is unable to regulate its activity and swims continuously (adapted from Zhang and Sillar 2012; Zhang et al. 2015).

We have also explored the effects of Na^+ pump manipulation in the lumbar spinal cord of neonatal mice, using two methods for evoking locomotor activity. Traditionally, a combination of drugs (dopamine, NMDA, serotonin) is applied to induce a continuous locomotor rhythm (Figure 4A). Under these conditions, blockade of Na^+ pumps using ouabain causes the rhythm frequency to increase (Figure 4B). Conversely, raising the levels of intracellular sodium using monensin, which indirectly activates the Na^+ pump, causes the opposite effect (Figure 4C). Whilst this reveals the importance of the Na^+ pump for frequency control, it obviously cannot address the role of Na^+ pumps in regulating intervals between locomotor episodes.

In order to address this question in a similar way to our earlier tadpole experiments, we switched to using dorsal root sensory stimulation to evoke individual, more natural bouts of locomotor activity (Figure 4D,E). In much the same way as in tadpoles, episode 2 is clearly influenced by episode 1 so long as the interval is shorter than 1 minute (Figure 4Ei). This relationship breaks down in the presence of ouabain such that episode 2 is now similar to episode 1 in duration, frequency and amplitude (Figure 4Eii).




Figure 4. Na^+ pump manipulation in the neonatal mouse preparation. Ai. Schematic depicting neonatal mouse spinal cord preparation. Aii. Glass suction electrodes are attached to the first or second lumbar ventral roots (L1, L2) on the left and right sides of an isolated spinal cord to record flexor-related activity, and a third electrode is attached to the fifth ventral root (L5) to record extensor-related activity. Aiii. Raw and rectified/integrated traces showing drug-induced activity on the left and right L2 roots and the right L5 root. B. Na^+ pump blockade increases the frequency of locomotor bursting. C.

Activation of the Na^+ pump has the opposite effect of slowing locomotor burst frequency. **D.** For sensory stimulation, an electrode was attached to the fourth or fifth dorsal root (L4 or L5) to deliver current pulses to initiate locomotion. **Ei.** When two episodes of locomotor output are evoked with a short interval (15 s), the second episode is both shorter and slower compared to this first episode. **Eii.** Following blockade of the Na^+ pump, not only are episodes longer and faster compared to control, but the interval relationship is abolished (Adapted from Picton et al. 2017).

Mechanism linking usAHP and A-current

The Na^+ pump-mediated usAHP clearly plays an important role in allowing locomotor networks to regulate their output in relation to past activity. However, it is not immediately obvious how the relatively modest membrane hyperpolarisation (~5 mV) caused by increased activation of the Na^+ pump can cause dramatic changes in neuronal excitability, especially since there is no obvious change in conductance. A likely possibility is that in different systems, different voltage-dependent currents are affected by the change in membrane potential. Two currents that appear to have important interactions with sodium pump currents in CPG networks are I_h and I_A (Kueh et al. 2016; Pulver and Griffith 2010; Zhang et al. 2015).

Pulver and Griffith (2010) showed in *Drosophila* larva motoneurons that the pump-mediated AHP brought the membrane potential into a range that caused the de-inactivation of an A-type potassium current, I_{shal} , which in turn introduced a delay to the first spike when activity resumed. Classically, channels mediating I_A are largely inactivated at the resting membrane potential but are de-inactivated by hyperpolarisation, so that when the neuron is next excited by a depolarising input the rate of depolarisation is slowed by I_A . We found precisely this mechanism at play in tadpole spinal neurons (Zhang et al. 2015). When a usAHP was induced by a high frequency train of action potentials (Figure 5A2), the delay to firing in response to a brief current pulse was longer compared with before the induction of a usAHP (Figure 5A3 vs. 5A1). The presence of a 4-AP-sensitive A-type potassium current was confirmed using voltage clamp recordings (Figure 5B). Whether a similar mechanism involving an A-type potassium current contributes to the role of the usAHP in neonatal mice is yet to be confirmed, but this possibility seems likely.




Figure 5. An A-type potassium current links the usAHP to inhibition of firing in *Xenopus* spinal neurons. **A.** Summary of the mechanism illustrating how the Na⁺ pump and A-type K⁺ current are involved in the short-term memory of motor network output. At rest, most A-type K⁺ channels are inactivated. Weak activity (1) does not increase Na⁺ pump current sufficiently to hyperpolarize the membrane potential so when the membrane potential is subsequently depolarized above threshold (2) most A-type K⁺ channels cannot be activated, and thus the first spike delay is unaffected. Stronger activity (2) can potentiate Na⁺ pump function and induce a larger pump current which hyperpolarizes the membrane potential (usAHP). This hyperpolarization removes the inactivation of A-type K⁺ channels, so that when depolarized above threshold (3), the A-type current is large enough to impede membrane depolarisation, prolonging first spike delay, and reducing the total number of spikes to a given depolarising input. **B.** Voltage clamp evidence for a 4-AP-sensitive A current in a spinal ascending interneuron (aIN). 4-AP preferentially blocks transient K⁺ currents. Red current trace is control, blue is in 4-AP and green is wash. Black trace is the difference in currents between control and 4-AP. (Adapted from Zhang et al. 2015).

Heterogenous distribution

The functional anatomy of the tadpole spinal network is known in considerable detail (Roberts et al. 2010), such that the presence or absence of a usAHP can be ascribed to each class of spinal neuron that participates in locomotory swimming. In three of the four main CPG classes (motoneurons (MNs), commissural interneurons (cINs) and ascending interneurons (aINs)), we found that approximately half of neurons display a usAHP; while in the other half of each subtype it is absent (Figure 6, Zhang and Sillar 2012). Furthermore, in one entire class, the excitatory rhythm-generating descending interneurons (dINs), the usAHP is absent altogether. The fact that dINs appear to be spared the influence of a usAHP presumably explains why some residual rhythm-generating capability remains regardless of how short the

inter-swim interval is (e.g. Figure 3B, 5s interval). However, the firing of dINs relies on rebound from mid-cycle inhibition coming from cINs on the contralateral side of the spinal cord, and therefore the impact of I_A on cIN firing will indirectly compromise dIN firing, and in turn the maintenance of the swim rhythm. The explanation for a lack of a pump current in dINs is yet to be determined, but one possibility is that they do not possess specific sodium pump isoforms responsible for mediating the usAHP (see discussion). Alternatively, the usAHP may be masked in this cell type by an equal, but opposite depolarising current, such as a persistent sodium current, or an I_h current, which may also become activated during intense spiking protocols (Darbon et al. 2004; Gullledge et al. 2013; Wang et al. 2012). This possibility is currently under investigation, with preliminary evidence suggesting that this may be the case.

A similar heterogenous usAHP distribution is present in the neonatal mouse CPG (Figure 6, Picton et al. 2017). For MNs, a very similar proportion to the tadpole (~40%) display the usAHP. For interneurons, there are many more classes in the mouse compared to the tadpole (Kiehn 2016), but around a quarter of unidentified interneurons that were recorded displayed a usAHP. This proportion is similar to that in tadpole interneurons when cINs, aINs and dINs are pooled. Although the identity of all the specific interneuron classes displaying a usAHP in neonatal mice is not yet known, one type of modulatory neurons, the cholinergic pitx2 class, was found to display a usAHP in around 60% of the population (Picton et al. 2017).




Figure 6. Heterogenous distribution of the usAHP among neuron types in *Xenopus* tadpoles (Zhang and Sillar 2012) and neonatal mice (Picton et al. 2017).

Discussion

Na⁺ pumps: intrinsic memory through a spike-rate monitor

Networks of neurons require the intrinsic capacity to monitor their own activity, allowing for the initiation of important homeostatic control mechanisms that adjust their output in light of past activity. Changes in neuronal and synaptic function often begin with changes in ionic conductances. The activity of a neuron may be reflected in changes in intracellular calcium concentration, leading to the activation of a range of downstream signalling pathways including protein phosphorylation and ion channel modulation. However, the clearance of calcium itself, mediated primarily by the calcium pump, is often relatively rapid (Benham et al. 1992), and therefore calcium influx is usually not considered to be responsible for electrical changes in the time scale of tens of seconds. Another ion intrinsically linked to neuronal activity is sodium, whose intracellular levels also rise rapidly during spiking before decaying slowly over tens of seconds after activity has ceased (Rose 2002). The Na⁺ pump is the primary means of restoring intracellular sodium concentrations. It is therefore strategically positioned both to homeostatically control changes in intracellular sodium levels resulting from neuronal firing, and to link neuronal activity to intrinsic excitability. It was shown as early as the 1950's that rises in intracellular sodium can cause a prolonged membrane hyperpolarisation (Coombs et al. 1955), and that this effect is mediated by the activation of the Na⁺ pump (Connelly 1959; Ritchie and Straub 1957).

This phenomenon has since been reported in a range of neuronal types at every level of the motor pathway. For example, pump-mediated AHPs have been reported in the *sensory neurons* of a range of species including insects (French 1989), lamprey (Parker et al. 1996), leech (Arganda et al. 2007; Baylor and Nicholls 1969; Scuri et al. 2002), crayfish (Nakajima and Takahashi 1966; Sokolove and Cooke 1971), frogs (Davidoff and Hackman 1980; Kobayashi et al. 1997), horseshoe crabs (Smith et al. 1968) and rats (Gordon et al. 1990). Similar post-tetanic AHP mechanisms mediated by the sodium pump have also been found in the *interneurons* of numerous species including the leech (Tobin and Calabrese 2005), *Aplysia* (Gage and Hubbard 1968; Pinsker and Kandel 1969) and rats (Darbon et al. 2002; 2003; Krey et al. 2010; Tsuzawa et al. 2015). Finally, the *motoneurons* of diverse species have also been shown to display a spike-dependent, pump-mediated hyperpolarisation, including in the motor axons of lizards (Morita et al. 1993), guinea pigs (del Negro et al. 1999), rats (Ballerini et al. 1997; Gage and Hubbard 1966) and humans (Kiernan et al. 2004; Vagg et al. 1998). In several networks, these activity-dependent hyperpolarisations have been shown to perform important roles in shaping the rhythmic output of the network itself; from neurosecretory networks in the snail brain (Nikolić et al. 2008, 2012; Tsai and Chen 1995), to rhythmic networks in the rat brain including the suprachiasmatic nucleus (Wang et al. 2004, 2006, 2012) and midbrain dopaminergic neurons (Johnson et al. 1992). More recently, sodium pumps have also been found to play an important role in shaping the output of hippocampal neurons (Azarias et al. 2013; Gullledge et al. 2013; Gustafsson and Wigström, 1983), striatal neurons (Azarias et al. 2013), cerebellar purkinje fibres (Forrest et al. 2012) and neurons in the auditory pathway (Kim et al. 2007, 2012). Sodium pumps thus play important roles throughout the

nervous system and across diverse species, and participate at every level of the motor pathway; from modifying sensory information, to the integration and relay of this information by interneuronal networks, right through to the regulation of the final motor output by motoneurons. However, only recently has the functional importance of the sodium pump as a spike-rate monitor been explored in depth in the spinal CPG networks controlling vertebrate locomotion.

Because of the close link between intracellular sodium levels and Na^+ pump activity, pharmacological tools that raise the levels of sodium in a neuron can be useful for studying the effects of increased Na^+ pump activity. Hence monensin, a sodium ionophore that exchanges one sodium ion intracellularly for one proton extracellularly, has been used extensively in studying sodium pumps (e.g. Kueh et al. 2016; Wang et al. 2012; Zhang et al. 2015). Monensin essentially acts as a proxy for intense spiking, imposing on neurons the pharmacological equivalent of a long train of high frequency action potentials. In both *Xenopus* and mouse spinal neurons, monensin increases Na^+ pump activity, hyperpolarising the membrane potential to the level attained by the usAHP. Locomotor activity, again in both species, becomes shorter and slower under monensin as if the network has been intensely active for a long period of time. Thus, monensin appears to chronically fatigue spinal networks by maximally activating the Na^+ pump autoregulation mechanism. Monensin has also recently been used to study the role of Na^+ pumps in the heartbeat network of the leech, where a fascinating interaction between a pump current and a depolarising I_h current was revealed (Kueh et al. 2016). Directly increasing intracellular sodium concentration using a modified intracellular solution could be used in future studies to confirm these findings.

Molecular and cellular basis for activity-dependent pump activation

The pump-based mechanisms that link future to past network activity transcend major phylogenetic boundaries and occur on multiple levels; from the molecular to the cellular and circuit levels.

At the molecular level, there is an emerging hypothesis that there exist both tonic and dynamic contributions of the sodium pump to membrane potential, and that these contributions rely partly on the heterogeneity of subunit composition of the pumps. In neurons in general, the α -subunit of the Na^+ pump takes one of two forms with different affinities for intracellular sodium; $\alpha 1$ (high affinity) or $\alpha 3$ (low affinity). Thus, at typical resting intracellular sodium levels, the $\alpha 1$ is maximally active, whilst the $\alpha 3$ remains inactive, or sub-maximally active, allowing it to act as a sensor for activity-dependent rises in sodium (Azarias et al. 2013; Dobretsov and Stimers, 2005). The subsequent increase in the activity of $\alpha 3$ -containing sodium pumps is thought to be responsible for generating the transient membrane hyperpolarisation that reduces the excitability of the neuron for tens of seconds. The different isoforms also have differential sensitivity to ouabain, such that low concentrations of ouabain, including those used in our experiments (1-3 μM), selectively block the $\alpha 3$ isoform (Blanco and Mercer 1998; Dobretsov and Stimers 2005). Our pharmacological experiments showing that the usAHP is blocked by these low concentrations of ouabain are therefore in support of the above hypothesis.

In mice, both $\alpha 1$ and $\alpha 3$ expression is found throughout the ventral and dorsal horns of the spinal cord, although $\alpha 3$ expression is more widespread (Edwards et al. 2013; Hieber et al. 1991; Watts et al. 1991). However, both $\alpha 1$ and $\alpha 3$ expression appears to be restricted to some neurons and not others. For instance, alpha-motoneurons predominantly express $\alpha 3$, whilst gamma-motoneurons predominantly express $\alpha 1$ (Edwards et al. 2013). The functional importance of this difference is not yet clear. Expression of $\alpha 3$ is also found in interneurons, and in our experiments, we specifically focused on $\alpha 3$ expression in one interneuron type, the cholinergic pitx2 cells (Zagoraïou et al. 2009). We found $\alpha 3$ expression in around half of this population, which broadly matches the number of pitx2 neurons found to display the usAHP (Picton et al. 2017). This is also similar to previous studies in rats which documented an activity-dependent, pump-mediated hyperpolarisation in around half of cultured spinal interneurons (Darbon et al. 2002, 2003). It will be important in future studies to further characterise $\alpha 3$ expression in other interneuron types. It will also be important to characterise developmental changes in $\alpha 3$ expression. For example, Calyx of Held neurons in young rats have lower expression of $\alpha 3$ compared to adults, and this is accompanied by a significantly smaller and shorter duration usAHP (Kim et al. 2007).

At the cellular level, we have partially characterised the details of the cascade of events in *Xenopus* tadpoles that link spinal neuron firing to network regulation. This cascade involves the spike-dependent accumulation of sodium ions, which in turn triggers an increase in ion exchange by the Na^+ pump, hyperpolarising the neuron. This hyperpolarisation de-inactivates an A-type potassium channel, and enhanced A-current delays spiking in a subset of spinal motor and interneurons when activity resumes, causing a collapse of swim network activity. Thus, swimming activity evoked within a minute after the end of previous swimming is both shorter in duration and slower in frequency, in a time-dependent manner. In mice, a similar physiological mechanism appears to be at play, but unsurprisingly, additional mechanisms of locomotor bout termination are likely to be involved. For example, unlike tadpoles, blockade of the Na^+ pump does not produce continuous locomotion, but merely extends the duration of evoked locomotor bouts (Picton et al. 2017). It is likely that synaptic depression plays a role in locomotor bout termination, a possibility that has been explored previously in rat spinal neurons in the context of the sodium pumps (Darbon et al. 2002, 2003; Rozzo et al. 2002). We also do not yet know whether A-currents play a role in neonatal mice. As we come to understand more about Na^+ pump currents, we will likely uncover species-specific mechanisms involving a range of other currents, such as the I_h current, which has been shown to have important interactions with pump currents in a number of different brain areas (Gulledge et al. 2013; Kim and von Gersdorff 2012; Rozzo et al. 2002; Trotier and Døving 1996).

Heterogeneity allied to circuit role

The usAHP is a powerful way of reducing network excitability. However, if it were to be homogenously expressed in all CPG neurons then there would be a distinct possibility that the network could render itself completely unresponsive. This, in turn,

could be catastrophic because of the requirement to retain a residual capacity to respond to potentially life-threatening stimuli such as an approaching predator. In both tadpole and neonatal mouse spinal locomotor networks there is strong evidence for a heterogeneous distribution of the usAHP among spinal CPG network components.

There are a number of possible explanations for the heterogeneous distribution of the usAHP among neuron subtypes in the spinal cord. One possibility, for which we have preliminary evidence in the mouse (described above) is that the ability of the pump to respond dynamically to intense activity requires the presence of an $\alpha 3$ -containing sodium pump, which is only recruited by high intracellular sodium concentrations achieved following intense neuronal firing. Alternatively, the α subunit may also be subject to direct phosphorylation in some neurons, but not others (Therien and Blostein 2000), which can tune the affinity of the subunit for sodium. A similar mechanism could also involve a set of accessory proteins, known as FXYD proteins, which are also subject to phosphorylation (Geering 2006). Thus, it will be important in future studies not only to establish the distribution of $\alpha 1$ and $\alpha 3$ subunit isoforms, but also the expression of FXYD proteins in the spinal cord.

The importance of the Na^+ pump as an intrinsic locomotor memory mechanism, and its high conservation through evolution, make it a useful target for a range of neuromodulators, and this could also explain differences in usAHP expression. The range of neuromodulators known to impinge on the Na^+ pump is extensive (Therien and Blostein 2000), but dopamine, serotonin and nitric oxide seem particularly important, especially in the spinal cord. Indeed, in mice we showed that the effects of Na^+ pump manipulation were dopamine-dependent, and that dopamine extends the duration of the usAHP (Picton et al. 2017). Whether this involves direct phosphorylation of sodium pumps, or via FXYD accessory proteins, or both, is a topic for future experiments.

Phylogenetic conservation

In this paper, we have reviewed the evidence that the activity-dependent increase in Na^+ pump activity, manifest as the usAHP, functions as a simple form of short-term motor memory in animals as diverse as fruitflies, frog tadpoles and neonatal mice. Modern amphibians and mammals diverged from a common ancestor that existed around 360 million years ago. The nervous system underwent dramatic changes to accommodate changes in lifestyle, morphology, and behavioural repertoire, with the number of neurons increasing from around 16 million in adult frogs to around 70 million in adult mice. However, many components of the nervous system are known to be highly conserved (Katz 2016; Katz and Harris-Warrick 1999; Keifer and Summers 2016). The basic architecture of many neural circuits appears to have been retained through evolutionary time, with extant species displaying variations on a theme rather than completely new circuit architecture. Thus, we can often identify conserved principles of circuit function and this often appears to be true for the circuits controlling locomotor behaviours, including at the cellular and molecular levels (Goulding and Pfaff 2005). The neuronal Na^+ pump is especially highly conserved between vertebrates in terms of its structure and function, with around 96%

cross-species similarity (Dobretsov and Stimers 2005; Takeyasu et al. 1990). This implies that the Na⁺ pump plays an important and conserved neuronal function. Our own mammalian lineage diverged from the common ancestor with mice around 65 million years ago (O'Leary et al. 2013), and so it will be interesting in future studies, especially with a rise in the use of human induced pluripotent stem cells (iPSCs), to study whether the sodium pumps embedded in human spinal motoneurons and interneurons also play a similar role in neuronal self-regulation.

Dysfunction of the Na⁺ pump

Na⁺ pumps are receiving increasing attention in mammalian systems not only for their importance for normal network function, but also for their relevance to both the ageing process and a range of debilitating diseases of the nervous system (de Lores Arnaiz and Ordieres 2014; Holm and Lykke-Hartmann 2016). The $\alpha 3$ Na⁺ pump isoform is highly expressed in the human brain and spinal cord (Peng et al. 1992) and several mutations in the gene encoding this subunit (*ATP1A3*) are known to cause at least three neurological disorders: Alternating Hemiplegia of Childhood (AHC, (Heinzen et al. 2012; Rosewich et al. 2012)); Rapid-onset Dystonia Parkinsonism (RDP, De Carvalho Aguiar et al. 2004; Rodacker et al. 2006); and Cerebellar ataxia, Areflexia, Pes cavus, Optic atrophy and Sensorineural hearing loss (CAPOS) syndrome (Demos et al. 2014). Furthermore, a wide range of other disorders are also known to involve changes in the activity of the $\alpha 3$ Na⁺ pump isoform. In recent studies, the $\alpha 3$ isoform has been shown to directly interact with both SOD1 (Martin et al. 2007; Ruegsegger et al. 2016), and α -synuclein (Shrivastava et al. 2015), in ALS and Parkinson's Disease mouse models, respectively. This aggregation leads to reduced $\alpha 3$ activity and a general inability to respond to rises in intracellular sodium (Ellis et al. 2003; Shrivastava et al. 2015). Given that dysfunction of $\alpha 3$ also contributes to epilepsy (Krishnan et al. 2015) and bipolar disorder (Kirshenbaum et al. 2012), it is possible that the inability to respond dynamically and homeostatically to activity-induced rises in intracellular sodium may be a general feature of pump disorders involving the $\alpha 3$ isoform (Azarias et al. 2013; Benarroch 2011).

Genetically modified zebrafish and rodent disease models have been used to explore the underlying mechanisms of Na⁺ pump deficiency. *ATP1A3* knockdown zebrafish display abnormal motor activity accompanied by depolarization of spinal sensory neurons (Doganli et al. 2013). Homozygous knock-out mice for $\alpha 1$ are embryonic lethal (James et al. 1999), whilst homozygous $\alpha 3$ knock-out mice die shortly after birth (Moseley et al. 2007). However, a number of $\alpha 3$ knock-in mouse lines have been developed and heterozygote mice all show severe motor deficits (DeAndrade et al. 2011; Hunanyan et al. 2015; Ikeda et al. 2013; Kirshenbaum et al. 2011; Moseley et al. 2007; Sugimoto et al. 2014). The hyperactivity phenotype in these mice is especially pronounced, with mutant mice showing almost continuous, high frequency locomotor activity compared to control mice. The $\alpha 3$ -mutation affects Na⁺ pumps throughout the nervous system, including presumably the spinal cord, and therefore this phenotype may relate to the role of the $\alpha 3$ Na⁺ pumps explored in this review. Indeed, this behavioural phenotype would be predicted by the effects

covered in this review using low concentrations of ouabain; namely, longer duration bouts of locomotion with a higher frequency of limb movements, and a general inability to regulate locomotion.

Summary

Na⁺/K⁺ exchange pumps are ubiquitously distributed, abundantly expressed and phylogenetically conserved proteins that are often viewed as molecular automata engaged exclusively in the maintenance of ionic distributions across cell membranes. Here, we have discussed recent data in *Xenopus* tadpoles, neonatal mice and also *Drosophila*, showing that Na⁺ pumps respond dynamically to changes in intracellular sodium that accompany intense neuronal firing. This capacity endows networks of the spinal cord with a homeostatic control mechanism to shape motor output in an activity-dependent manner. Moreover, despite the ubiquity of Na⁺ pump distribution among network neurons, their ability to respond homeostatically to the changes in intracellular sodium triggered by activity may result from the highly targeted insertion of $\alpha 3$ -containing pumps in selected neurons and neuronal subtypes. The possibility that the balance of $\alpha 1$ to $\alpha 3$ expression is a mutable entity that can change during development, or with circuit use, is an exciting idea that should be pursued in the future.

References

- Arganda S, Guantes R, de Polavieja GG. Sodium pumps adapt spike bursting to stimulus statistics. *Nat Neurosci* 11: 1467–1473, 2007.
- Azarias G, Kruusmägi M, Connor S, Akkuratov EE, Liu XL, Lyons D, Brismar H, Broberger C, Aperia A. A specific and essential role for Na,K-ATPase $\alpha 3$ in neurons co-expressing $\alpha 1$ and $\alpha 3$. *J Biol Chem* 288: 2734–2743, 2013.
- Ballerini L, Bracci E, Nistri A. Pharmacological block of the electrogenic sodium pump disrupts rhythmic bursting induced by strychnine and bicuculline in the neonatal rat spinal cord. *J Neurophysiol* 77: 17–23, 1997.
- Baylor DA, Nicholls JG. After-effects of nerve impulses on signalling in the central nervous system of the leech. *J Physiol* 203: 571–589, 1969.
- Benarroch EE. Na⁺,K⁺-ATPase: Functions in the nervous system and involvement in neurologic disease. *Neurology* 76: 287–293, 2011.
- Benham CD, Evans ML, McBain CJ. Ca²⁺ efflux mechanisms following depolarization evoked calcium transients in cultured rat sensory neurones. *J Physiol* 455: 567–583, 1992.
- Blanco G, Mercer RW. Isozymes of the Na-K-ATPase: heterogeneity in structure, diversity in function. *Am J Physiol* 275: 633–650, 1998.
- Carpenter DO, Alving BO. A contribution of an electrogenic Na⁺ pump to membrane potential in *Aplysia* neurons. *J Gen Physiol* 52: 1–21, 1968.
- Connelly CM. Recovery Processes and Metabolism of Nerve. *Rev Mod Phys* 31:475–484, 1959.
- Coombs JS, Eccles JC, Fatt P. The electrical properties of the motoneurone

557 membrane. *J physiol* 130: 291–325, 1955.

558 Darbon P, Scicluna L, Tscherter A, Streit J. Mechanisms controlling bursting activity
559 induced by disinhibition in spinal cord networks. *Eur J Neurosci* 15: 671–683,
560 2002.

561 Darbon P, Tscherter A, Yvon C, Streit J. Role of the electrogenic Na/K pump in
562 disinhibition-induced bursting in cultured spinal networks. *J neurophysiol* 90:
563 3119–3129, 2003.

564 Darbon P, Yvon C, Legrand JC, Streit J. I_{NaP} underlies intrinsic spiking and rhythm
565 generation in networks of cultured rat spinal neurons. *Eur J Neurosci* 20: 976–
566 988, 2004.

567 Davidoff RA, Hackman JC. Hyperpolarization of frog primary afferent fibres caused
568 by activation of a sodium pump. *J Physiol* 302: 297–309, 1980.

569 del Negro CA, Hsiao C-F, Chandler SH. Outward currents influencing bursting
570 dynamics in guinea pig trigeminal motoneurons. *J Neurophysiol* 81: 1478–1485,
571 1999.

572 De Andrade MP, Yokoi F, van Groen T, Lingrel JB, Li Y. Characterisation of ATP1 α 3
573 mutant mice as a model of rapid-onset dystonia with parkinsonism. *Behav Brain*
574 *Res* 216: 659-665, 2011.

575 de Carvalho Aguiar P, Sweadner KJ, Penniston JT, Zaremba J, Liu L, Caton M,
576 Linazasoro G, Borg M, Tijssen MA, Bressman SB, Dobyns WB, Brashear A,
577 Ozelius LJ. Mutations in the Na⁺/K⁺-ATPase α 3 gene *ATP1A3* are associated
578 with rapid-onset dystonia parkinsonism. *Neuron* 43: 169–175, 2004.

579 de Lores Arnaiz GR, Ordieres MGL. Brain Na⁺, K⁺-ATPase activity in aging and
580 disease. *Int J Biomed Sci* 10: 85–102, 2014.

581 Demos MK, van Karnebeek CD, Ross CJ, Adam S, Shen Y, Zhan SH, Shyr C,
582 Horvath G, Suri M, Fryer A, Jones SJ, Friedman JM, FORGE Canada
583 Consortium. A novel recurrent mutation in ATP1A3 causes CAPOS syndrome.
584 *Orphanet J Rare Dis* 9: 15, 2014.

585 Dobretsov M, Stimers JR. Neuronal function and alpha3 isoform of the Na/K-ATPase.
586 *Front Biosci* 10: 2373–2396, 2005.

587 Doganli C, Beck HC, Ribera AB, Oxvig C, Lykke-Hartmann K. α 3 Na⁺/K⁺-ATPase
588 deficiency causes brain ventricle dilation and abrupt embryonic motility in
589 zebrafish. *J Biol Chem* 288: 8862–8874, 2013.

590 Edwards IJ, Bruce G, Lawrenson C, Howe L, Clapcote SJ, Deuchars SA, Deuchars J.
591 Na⁺/K⁺ ATPase α 1 and α 3 isoforms are differentially expressed in α - and γ -
592 motoneurons. *J Neurosci* 33: 9913–9919, 2013.

593 Ellis DZ, Rabe J, Sweadner KJ. Global loss of Na,K-ATPase and its nitric oxide-
594 mediated regulation in a transgenic mouse model of amyotrophic lateral
595 sclerosis. *J Neurosci* 23: 43–51, 2003.

596 Engl E, Attwell D. Non-signalling energy use in the brain. *J Physiol* 593: 3417–3429,
597 2015.

598 Forrest MD, Wall MJ, Press DA, Feng J. The sodium-potassium pump controls the
599 intrinsic firing of the cerebellar purkinje neuron. *PLoS ONE* 7: e51169, 2012.

600 French AS. Ouabain selectively affects the slow component of sensory adaptation in
601 an insect mechanoreceptor. *Brain Res* 504: 112–114, 1989.

602 Gage PW, Hubbard JI. The origin of the post-tetanic hyperpolarization of mammalian
603 motor nerve terminals. *J Physiol* 184: 335–352, 1966.

604 Gandevia SC. Spinal and supraspinal factors in human muscle fatigue. *Physiol Rev*
605 81: 1725–1789, 2001.

606 Geering K. FXYD proteins: new regulators of Na-K-ATPase. *Am J Physiol Renal*
607 *Physiol* 290: 241–250, 2006.

608 Gordon TR, Kocsis JD, Waxman SG. Electrogenic pump (Na⁺/K⁺-ATPase) activity in
609 rat optic nerve. *Neuroscience* 37: 829–837, 1990.

610 Goulding M, Pfaff SL. Development of circuits that generate simple rhythmic
611 behaviors in vertebrates. *Curr Opin Neurobiol* 15: 14–20, 2005.

612 Gullledge AT, Dasari S, Onoue K, Stephens EK, Hasse JM, Avesar D. A sodium-
613 pump-mediated afterhyperpolarization in pyramidal neurons. *J Neurosci* 33:
614 13025–13041, 2013.

615 Gustafsson B, Wigström H. Hyperpolarization following long-lasting tetanic activation
616 of hippocampal pyramidal cells. *Brain Res* 275: 159–163, 1983.

617 Harris-Warrick RM, Marder E. Modulation of neural networks for behavior. *Annu Rev*
618 *Neurosci* 14: 39–57, 1991.

619 Heinzen EL, Swoboda KJ, Hitomi Y, Gurrieri F, Nicole S, de Vries B, Tiziano FD,
620 Fontaine B, Walley NM, Heavin S, Panagiotakaki E; European Alternating
621 Hemiplegia of Childhood (AHC) Genetics Consortium.; Biobanca e Registro
622 Clinico per l'Emiplegia Alternante (I.B.AHC) Consortium.; European Network for
623 Research on Alternating Hemiplegia (ENRAH) for Small and Medium-sized
624 Enterprise (SMEs) Consortium., Fiori S, Abiusi E, Di Pietro L, Sweney MT,
625 Newcomb TM, Viollet L, Huff C, Jorde LB, Reyna SP, Murphy KJ, Shianna KV,
626 Gumbs CE, Little L, Silver K, Ptáček LJ, Haan J, Ferrari MD, Bye AM, Herkes
627 GK, Whitelaw CM, Webb D, Lynch BJ, Uldall P, King MD, Scheffer IE, Neri G,
628 Arzimanoglou A, van den Maagdenberg AM, Sisodiya SM, Mikati MA, Goldstein
629 DB. De novo mutations in ATP1A3 cause alternating hemiplegia of childhood.
630 *Nat Genet* 44: 1030–1034, 2012.

631 Hieber V, Siegel GJ, Fink DJ, Beaty MW, Mata M. Differential distribution of (Na, K)-
632 ATPase α isoforms in the central nervous system. *Cell Mol Neurobiol*, 11: 253–
633 262, 1991.

634 Hodgkin AL, Keynes RD. Experiments on the injection of substances into squid giant
635 axons by means of a microsyringe. *J Physiol* 131: 592–616, 1956.

636 Holm TH, Lykke-Hartmann K. Insights into the pathology of the $\alpha 3$ Na⁺/K⁺-ATPase in
637 neurological disorders; lessons from animal models. *Front Physiol* 7: 209, 2016.

638 Hunanyan AS, Fainberg NA, Linabarger M, Arehart E, Leonard AS, Adil SM, Helseth
639 AR, Swearingen AK, Forbes SL, Rodriguiz RM, Rhodes T, Yao X, Kibbi N,

640 Hochman DW, Wetzel WC, Hochgeschwender U, Mikati MA. Knock-in mouse
641 model of alternating hemiplegia of childhood: behavioral and electrophysiologic
642 characterization. *Epilepsia* 56: 82–93, 2015.

643 Ikeda K, Satake S, Onaka T, Sugimoto H, Takeda N, Imoto K, Kawakami K.
644 Enhanced inhibitory neurotransmission in the cerebellar cortex of Atp1α3-
645 deficient heterozygous mice. *J physiol* 591: 3433–49, 2013.

646 James PF, Grupp IL, Grupp G, Woo AL, Askew GR, Croyle ML, Walsh RA, Lingrel
647 JB. Identification of a specific role for the Na,K-ATPase α2 isoform as a
648 regulator of calcium in the heart. *Mol Cell* 3: 555–563, 1999.

649 Johnson SW, Seutin V, North R. Burst firing in dopamine neurons induced by N-
650 methyl-D-aspartate: role of electrogenic sodium pump. *Science* 258: 665–667,
651 1992.

652 Katz PS. Evolution of central pattern generators and rhythmic behaviours. *Philos*
653 *trans R Soc Lond B Biol Sci* 371: 20150057, 2016.

654 Katz PS, Harris-Warrick MR. The evolution of neural circuits underlying species-
655 specific behavior. *Curr Opin Neurobiol* 9: 628–633, 1999.

656 Keifer J, Summers CH. Putting the “biology” back into “neurobiology”: the strength of
657 diversity in animal model systems for neuroscience research. *Front Syst*
658 *Neurosci* 10: 1–9, 2016.

659 Kiehn O. Decoding the organization of spinal circuits that control locomotion. *Nat*
660 *Rev Neurosci* 17: 224–238, 2016.

661 Kiernan MC, Lin CS-Y, Burke D. Differences in activity-dependent hyperpolarization
662 in human sensory and motor axons. *J Physiol* 558: 341–349, 2004.

663 Kim JH, Sizov I, Dobretsov M, von Gersdorff H. Presynaptic Ca²⁺ buffers control the
664 strength of a fast post-tetanic hyperpolarization mediated by the α3 Na⁺/K⁺-
665 ATPase. *Nat Neurosci* 10: 196–205, 2007.

666 Kim JH, von Gersdorff H. Suppression of spikes during posttetanic hyperpolarization
667 in auditory neurons: the role of temperature, I_h currents, and the Na⁺-K⁺-ATPase
668 pump. *J Neurophysiol* 108: 1924–1932, 2012.

669 Kirshenbaum GS, Saltzman K, Rose B, Petersen J, Vilsen B, Roder JC. Decreased
670 neuronal Na⁺,K⁺-ATPase activity in Atp1α3 heterozygous mice increases
671 susceptibility to depression-like endophenotypes by chronic variable stress.
672 *Genes Brain Behav* 10: 542–550, 2011.

673 Kirshenbaum GS, Clapcote SJ, Duffy S, Burgess CR, Petersen J, Jarowek KJ, Yücel
674 YH, Cortez MA, Snead OC 3rd, Vilsen B, Peever JH, Ralph MR, Roder JC.
675 Mania-like behavior induced by genetic dysfunction of the neuron-specific
676 Na⁺,K⁺-ATPase α3 sodium pump. *Proc Natl Acad Sci USA* 108: 18144–18149,
677 2012.

678 Kobayashi JI, Ohta M, Terada Y. Evidence for the involvement of Na⁺-K⁺ pump and
679 K⁺ conductance in the post-tetanic hyperpolarization of the tetrodotoxin-resistant
680 C-fibers in the isolated bullfrog sciatic nerve. *Neurosci Lett* 236: 171–174, 1997.

681 Krey RA, Goodreau AM, Arnold TB, Del Negro CA. Outward currents contributing to

682 inspiratory burst termination in preBötzinger complex neurons of neonatal mice
 683 studied *in vitro*. *Front Neural Circuits* 4: 124, 2010.

684 Krishnan AV, Lin CS, Kiernan MC. Activity-dependent excitability changes suggest
 685 Na⁺/K⁺ pump dysfunction in diabetic neuropathy. *Brain* 131: 1209–1216, 2008.

686 Krishnan GP, Filatov G, Shilnikov A, Bazhenov M. Electrogenic properties of the
 687 Na⁺/K⁺ ATPase controls transitions between normal and pathological brain
 688 states. *J Neurophysiol* 113: 3356–3374, 2015.

689 Kueh D, Barnett WH, Cymbalyuk GS, Calabrese RL. Na⁺/K⁺ pump interacts with the
 690 *h*-current to control bursting activity in central pattern generator neurons of
 691 leeches. *eLife* 5: e19322, 2016.

692 Martin LJ, Liu Z, Chen K, Price AC, Pan Y, Swaby JA, Golden WC. Motor neuron
 693 degeneration in amyotrophic lateral sclerosis mutant superoxide dismutase-1
 694 transgenic mice: mechanisms of mitochondriopathy and cell death. *J Comp*
 695 *Neurol* 500: 20–46, 2007.

696 Morita K, David G, Barrett JN, Barrett EF. Posttetanic hyperpolarization produced by
 697 electrogenic Na⁺-K⁺ pump in lizard axons impaled near their motor terminals. *J*
 698 *Neurophysiol* 70: 1874–1884, 1993.

699 Moseley AE, Williams MT, Schaefer TL, Bohanan CS, Neumann JC, Behbehani MM,
 700 Vorhees CV, Lingrel JB. Deficiency in Na,K-ATPase α isoform genes alters
 701 spatial learning, motor activity, and anxiety in mice. *J Neurosci* 27: 616–626,
 702 2007.

703 Nakajima S, Takahashi K. Post-tetanic hyperpolarization and electrogenic Na pump
 704 in stretch receptor neurone of crayfish. *J Physiol* 187: 105–127, 1966.

705 Nikolić L, Kartelija G, Nedeljković M. Effect of static magnetic fields on bioelectric
 706 properties of the Br and N1 neurons of snail *Helix pomatia*. *Comp Biochem Phys*
 707 *A* 151: 657–663, 2008.

708 Nikolić L, Todorović N, Zakrzewska J, Stanić M, Rauš S, Kalauzi A, Janać B.
 709 Involvement of Na⁺/K⁺ pump in fine modulation of bursting activity of the snail Br
 710 neuron by 10 mT static magnetic field. *J Comp Physiol A Neuroethol Sens*
 711 *Neural Behav Physiol* 198: 525–540, 2012.

712 O’Leary MA, Bloch JI, Flynn JJ, Gaudin TJ, Giallombardo A, Giannini NP, Goldberg
 713 SL, Kraatz BP, Luo ZX, Meng J, Ni X, Novacek MJ, Perini FA, Randall ZS,
 714 Rougier GW, Sargis EJ, Silcox MT, Simmons NB, Spaulding M, Velazco PM,
 715 Weksler M, Wible JR, Cirranello AL. The placental mammal ancestor and the
 716 post-K-Pg radiation of placentals. *Science* 339: 662–667, 2013.

717 Parker D, Hill R, Grillner S. Electrogenic pump and a Ca(2⁺)-dependent K⁺
 718 conductance contribute to a posttetanic hyperpolarization in lamprey sensory
 719 neurons. *J Neurophysiol* 76: 540–553, 1996.

720 Picton LD, Nascimento F, Broadhead MJ, Sillar KT, Miles GB. Sodium pumps
 721 mediate activity-dependent changes in mammalian motor networks. *J Neurosci*,
 722 37: 906–921, 2017.

723 Peng JHF, Zeng Y, Parker Jr JC. Highly ouabain-sensitive Alpha3 isoform of Na⁺,K⁺-
 724 ATPase in human brain. *J Neurochem* 58, 1180–1183, 1992.

- 725 Pinsker H, Kandel ER. Synaptic activation of an electrogenic sodium pump. *Science*
726 163: 931-935, 1969.
- 727 Pulver SR, Griffith LC. Spike integration and cellular memory in a rhythmic network
728 from Na^+/K^+ pump current dynamics. *Nat neurosci* 13: 53–59, 2010.
- 729 Ranieri F, Di Lazzaro V. The role of motor neuron drive in muscle fatigue.
730 *Neuromuscul Disord* 22 Suppl 3: S157–S161, 2012.
- 731 Reklung JC, Funk GD, Bayliss DA, Dong XW, Feldman JL. Synaptic control of
732 motoneuronal excitability. *Physiol Rev*, 80: 767–852, 2000.
- 733 Ritchie JM, Straub RW. The hyperpolarization which follows activity in mammalian
734 non-medullated fibres. *J Physiol* 136: 80–97, 1957.
- 735 Roberts A, Li W-C, Soffe SR. How neurons generate behavior in a hatchling
736 amphibian tadpole: an outline. *Front Behav neurosci* 4: 16, 2010.
- 737 Rodacker V, Toustrup-Jensen M, Vilsen B. Mutations Phe785Leu and Thr618Met in
738 Na^+/K^+ -ATPase, associated with familial rapid-onset dystonia parkinsonism,
739 interfere with Na^+ interaction by distinct mechanisms. *J Biol Chem* 281: 18539–
740 18548, 2006.
- 741 Rose CR. Na^+ signals at central synapses. *Neuroscientist* 8: 532–539, 2002.
- 742 Rosewich H, Thiele H, Ohlenbusch A, Maschke U, Altmüller J, Frommolt P, Zirn B,
743 Ebinger F, Siemes H, Nürnberg P, Brockmann K, Gärtner J. Heterozygous de-
744 novo mutations in ATP1A3 in patients with alternating hemiplegia of childhood: a
745 whole-exome sequencing gene-identification study. *Lancet Neurol* 11: 764–773,
746 2012.
- 747 Rossi A, Rossi S, Ginanneschi F. Activity-dependent changes in intrinsic excitability
748 of human spinal motoneurons produced by natural activity. *J Neurophysiol* 108:
749 2473–2480, 2012.
- 750 Rozzo A, Ballerini L, Abbate G, Nistri A. Experimental and modeling studies of novel
751 bursts induced by blocking Na^+ pump and synaptic inhibition in the rat spinal
752 cord. *J Neurophysiol* 88: 676–691, 2002.
- 753 Ruegsegger C, Maharjan N, Goswami A, Filézac de L'Etang A, Weis J, Troost D,
754 Heller M, Gut H, Saxena S. Aberrant association of misfolded SOD1 with Na^+/K^+
755 ATPase- $\alpha 3$ impairs its activity and contributes to motor neuron vulnerability in
756 ALS. *Acta Neuropathol* 131: 427-451, 2016.
- 757 Scuri R, Mozzachiodi R, Brunelli M. Activity-dependent increase of the AHP
758 amplitude in T sensory neurons of the leech. *J Neurophysiol* 88: 2490-2500,
759 2002.
- 760 Shrivastava AN, Redeker V, Fritz N, Pieri L, Almeida LG, Spolidoro M, Liebmann T,
761 Bousset L, Renner M, Léna C, Aperia A, Melki R, Triller A. α -synuclein
762 assemblies sequester neuronal $\alpha 3$ - Na^+/K^+ -ATPase and impair Na^+ gradient.
763 *EMBO J* 34: 2408–2423, 2015.
- 764 Sokolove PG, Cooke IM. Inhibition of impulse activity in a sensory neuron by an
765 electrogenic pump. *J Gen Physiol* 57: 125–163, 1971.

766 Smith TG, Stell WK, Brown JE, Freeman JA, Murray GC. A role for the sodium pump
767 in photoreception in *Limulus*. *Science* 162: 456–458, 1968.

768 Storm JF. Action potential repolarization and a fast after-hyperpolarization in rat
769 hippocampal pyramidal cells. *J Physiol* 385: 733–759, 1987.

770 Sugimoto H, Ikeda K, Kawakami K. Heterozygous mice deficient in Atp1α3 exhibit
771 motor deficits by chronic restraint stress. *Behav Brain Res* 272: 100–110, 2014.

772 Takeyasu K, Lemas V, Fambrough DM. Stability of Na⁺-K⁺-ATPase alpha-subunit
773 isoforms in evolution. *Am J Physiol* 259: 619–630, 1990.

774 Therien AG, Blostein R. Mechanisms of sodium pump regulation. *Am J Physiol Cell*
775 *Physiol* 279: 541–566, 2000.

776 Tobin A-E, Calabrese RL. Myomodulin increases Ih and inhibits the Na/K pump to
777 modulate bursting in leech heart interneurons. *J Neurophysiol* 94: 3938–3950,
778 2005.

779 Trotier D, Døving KB. Direct influence of the sodium pump on the membrane
780 potential of vomeronasal chemoreceptor neurones in frog. *J physiol* 490: 611–
781 621, 1996.

782 Tsai, MC, Chen, YH. Bursting firing of action potentials in central snail neurons
783 elicited by D-amphetamine: role of the electrogenic sodium pump. *Comp*
784 *Biochem Physiol C Pharmacol Toxicol Endocrinol* 111: 131–141, 1995.

785 Tsuzawa K, Yazawa I, Shakuo T, Ikeda K, Kawakami K, Onimaru H. Effects of
786 ouabain on respiratory rhythm generation in brainstem-spinal cord preparation
787 from newborn rats and in decerebrate and arterially perfused in situ preparation
788 from juvenile rats. *Neuroscience* 286, 404–411, 2015.

789 Vagg R, Mogyoros I, Kiernan MC, Burke D. Activity-dependent hyperpolarization of
790 human motor axons produced by natural activity. *J Physiol* 507: 919–925, 1998.

791 Wang H-Y, Huang R-C. Diurnal modulation of the Na⁺/K⁺-ATPase and spontaneous
792 firing in the rat retinorecipient clock neurons. *J Neurophysiol* 92: 2295–2301,
793 2004.

794 Wang Y-C, Yang J-J, Huang R-C. Intracellular Na⁺ and metabolic modulation of
795 Na/K pump and excitability in the rat suprachiasmatic nucleus neurons. *J*
796 *Neurophysiol* 108: 2024–2032, 2012.

797 Watts AG, Sanchez-Watts G, Emanuel JR, Levenson R. Cell-specific expression of
798 mRNAs encoding Na⁺,K⁺-ATPase α- and β-subunit isoforms within the rat
799 central nervous system. *Proc Natl Acad Sci U S A* 88: 7425–7429, 1991.

800 Zagoraiou L, Akay T, Martin JF, Brownstone RM, Jessell TM, Miles GB. A cluster of
801 cholinergic premotor interneurons modulates mouse locomotor activity. *Neuron*,
802 64: 645–662, 2009.


803 Zhang H-Y, Picton L, Li WC, Sillar KT. Mechanisms underlying the activity-
804 dependent regulation of locomotor network performance by the Na⁺ pump. *Sci*
805 *Rep* 5: 16188, 2015.

806 Zhang H-Y, Sillar KT. Short-term memory of motor network performance via activity-


807 dependent potentiation of Na^+/K^+ pump function. *Curr Biol* 22: 526–531, 2012.

808


Ai




Aii




Bi




Bii










Ci




Cii









Ai




Aii




Aiii




B




C




D





Ei Control




Eii Ouabain







Xenopus




Motoneurons



Interneurons



dINs




aINs




cINs

usAHP
No usAHP


Mouse



Motoneurons



Interneurons



PitX2



Other